

AD-A043 094

MICHIGAN UNIV ANN ARBOR DEPT OF PHYSICS  
RESONANT AND NON-RESONANT OPTICAL FREQUENCY MIXING IN SIMPLE MO--ETC(U)  
APR 77 J F WARD

F/G 20/5

AF-AFOSR-2302-72

UNCLASSIFIED


AFOSR-TR-77-0981

NL

1 of 1  
ADA043094



END  
DATE  
FILMED  
9-77  
DDC



• MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
18 AFOSR TR-77-0981			
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED	
RESONANT AND NON-RESONANT OPTICAL FREQUENCY MIXING IN SIMPLE MOLECULAR SYSTEMS.		Scientific - Final rept.	
6. PERFORMING ORG. REPORT NUMBER		8. CONTRACT OR GRANT NUMBER(s)	
10. AUTHOR(s) J. E. Ward		15. AF- AFOSR 22-2302-72	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics University of Michigan Ann Arbor MI 48109		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 9751-02 12 42 61102F	
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NP Bolling AFB, Bldg. 410 Washington DC 20332		12. REPORT DATE Apr 1977	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 8	
12 10p.		15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A technique based on dc-electric-field induced optical second-harmonic generation for measuring second- and third-order electric polarizabilities of molecules in the vapor phase has been developed. The technique has been used to produce detailed and precise data for a wide range of molecules which exist either as gases or liquids at ambient temperature. Hypolarizabilities give rise to phenomena such as optical frequency mixing, modulation and demodulation. Vapors offer technological advantages over crystals for these purposes but the main motivation in our present work is to gain a detailed, fundamental - ORCA			

ADA 043094

AU NO.

DDC FILE COPY

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

228 720 SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

understanding of these properties in the simplest possible molecular systems. This ruby laser radiation is in resonance with two-photon transition in cesium and this system is a useful one in which to study a number of resonant nonlinear optical processes: two-photon absorption and subsequent fluorescence, third harmonic generation, and four-photon parametric oscillation.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dis	SPECIAL
P	

UNCLASSIFIED



AFOSR-TR- 77- 0981

THE UNIVERSITY OF MICHIGAN  
ANN ARBOR

DEPARTMENT OF PHYSICS

PHYSICS-ASTRONOMY BUILDING  
TEL. NO. 313-764-3459

April 28, 1977

Approved for public release;  
distribution unlimited.

Lt. Col. Gordon Wepfer  
AFOSR/HP  
Bolling Air Force Base  
Washington, D.C. 20332

Dear Colonel Wepfer:

Final Report AFOSR 72-2302

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

Resonant and Non-Resonant Optical Frequency Mixing in Simple Molecular Systems.

I. MOLECULAR HYPERPOLARIZABILITIES (see Journal Articles 1, 2, 3, 5, 10, 11, 12)

We have established and developed a technique based on dc-electric-field induced optical second-harmonic generation for measuring second- and third-order electric polarizabilities of molecules in the vapor phase. The technique has been used to produce detailed and precise data for a wide range of molecules which exist either as gases or liquids at ambient temperature. The molecules studied include:

Inert gases: He, Ne, Ar, Kr, Xe

Thirteen Halogenated methanes:  $CX_nY_{4-n}$   $n=1-4$ ; X,Y=H, F, Cl and  $CBrF_3$

Miscellaneous molecules:  $H_2$ ,  $O_2$ ,  $N_2$ , CO, NO,  $H_2S$ ,  $(CH_3)_2O$ ,  $H_2O$ ,  $CO_2$ ,  $NH_3$ ,  
 $SF_6$ .

Hyperpolarizabilities give rise to phenomena such as optical frequency mixing, modulation and demodulation. Vapors offer technological advantages over crystals for these purposes but the main motivation in our present work is to gain a detailed, fundamental understanding of these properties in the simplest possible molecular systems.

For even these simple molecules, ab initio calculations are difficult, typically differing (in the few cases where a calculation has been done) from

experimental results by a factor three whereas the experimental uncertainties are typically better than 10%. Particularly exciting to us is the recent interest shown by Dr. Rod Bartlett of Battelle in improving the ab initio calculations and we look forward to a fruitful interaction here.

We have been interested in less fundamental theories which aim to predict nonlinear properties either from linear properties or from the nonlinear properties of related systems. (This is the only type of approach which has proved useful in predicting the nonlinear optical properties of crystals.) We have been particularly interested in the Bond Additivity Model (based on the notion that a C-Cl bond, for example, has the same nonlinear properties whether it is in a  $\text{CCl}_4$  molecule or a  $\text{CF}_3\text{Cl}$  molecule) and in modifications of that model which give useful results without entirely sacrificing simplicity. (See Journal Articles 10, 11, 12.)

Comparison of our vapor-phase measurements with related measurements on liquids is of particular interest since it serves as a probe of the molecular environment in the liquid state.

These topics are all discussed in the Journal Articles. Data to be published in Journal Article 12 is included here (see Appendix).

## II. RESONANT NONLINEAR OPTICAL PROCESSES (see Journal Articles 4, 7, 13)

A ruby laser serves as a narrow band source of limited but sufficient tuneability in the region of 6935Å. This radiation is in resonance with the two-photon transition  $6^2S_{1/2} - 9^2D_{3/2}$  in cesium and this system is a useful one in which to study a number of resonant nonlinear optical processes: two-photon absorption and subsequent fluorescence, third harmonic generation, and four-photon parametric oscillation.

Resonances offer the possibility of dramatically enhanced nonlinear optical processes. In addition to processes mentioned above, Raman processes, lasing,

optical Stark shifts, multiphoton ionization and distribution of the population over a large number of excited states all occur simultaneously and must be considered together to get a valid picture of the overall interaction. The spatial and temporal dependence of the laser beam and spectral hole burning must also be considered. An understanding of this situation will be fascinating from a fundamental point of view and invaluable technologically in making the best use of resonant enhancement in frequency conversion applications. Details of our present, far-from complete understanding are given in Journal Articles 4 and 7. Work continues on the preparation of Journal Article 13 and Mr. Smith's thesis, which is almost complete.

### III. OTHER TOPICS

a) DC-Electric-Field-Induced Optical Second-Harmonic Generation as a Probe of the Focal Region of a Laser Beam (see Journal Articles 6, 8).

Second-harmonic generation in this process depends on a spatial convolution of the dc and optical fields. Since the dc field geometry is known, harmonic generation can be used to yield information on the geometry of the laser beam focus. Air serves as the nonlinear medium and the technique gives detailed information which is difficult to obtain by any other method. Details are given in Journal Articles 6 and 8.

b) DC-Electric-Field-Induced Optical Rectification in Nitrobenzene (see Journal Article 9).

This process results in the generation of a voltage across a Kerr cell during the passage of a laser beam through it. It is theoretically interesting because of its relation to the Kerr effect and, perhaps, of practical interest as a basis for a fast (if insensitive) radiation detector. For details see Journal Article 9.



#### JOURNAL ARTICLES

1. "Nonlinear Optical Measurement of Dispersion in Gases," R.S. Finn and J.F. Ward, *Applied Optics* **21**, 11 (1972).
2. "Measurement of the Hyperpolarizability Ratio  $\chi_{yyyy}(-2\omega;0,\omega,\omega)/\chi_{yyxx}(-2\omega;0,\omega,\omega)$  for the Inert Gases," Irving J. Bigio and J.F. Ward, *Phys. Rev. A* **9**, 35 (1974).
3. "Measurements of Hyperpolarizabilities for Some Halogenated Methanes," R.S. Finn and J.F. Ward, *J. Chem. Phys.* **60**, 454 (1974).
4. "Two-photon resonant, Optical Third-harmonic Generation in Cesium Vapor," K.M. Leung, J.F. Ward, and B.J. Orr, *Phys. Rev. A* **9**, 2440-2448 (1974).
5. "Molecular Second- and Third-order Polarizabilities from Measurements of Second-harmonic Generation in Gases," J.F. Ward and Irving J. Bigio, *Phys. Rev. A* **11**, 60-66 (1975).
6. "Electric Field Induced Harmonic Generation as a Probe of the Focal Region of a Laser Beam," Irving J. Bigio, R.S. Finn, and J.F. Ward, *Applied Optics* **14**, 336-342 (1975).
7. "Saturation of Two-Photon-Resonant Processes in Cesium Vapor," J.F. Ward and A.V. Smith, *Phys. Rev. Letters* **35**, 653 (1975).
8. "Laser Beam Characteristics Using DC-Induced Second Harmonic Generation," J.K. Guha, *Journal of Applied Optics* **15**, 2381 (1976).
9. "Electric Field Induced Optical Rectification in Nitrobenzene," J.F. Ward and J.K. Guha, *Appl. Phys. Letts.* **30**, 276 (1977).
10. "Measurements of Nonlinear Optical Polarizabilities for Some Halogenated Methanes: The Role of Bond-Bond Interactions," C.K. Miller and J.F. Ward. (Submitted to *Physical Review*.)



11. "An Interacting Segment Model of Molecular Electric Tensor Properties: Theory and Application to Electric Dipole Moments of the Halogenated Methanes," C.K. Miller, B.J. Orr, and J.F. Ward. (Submitted to J. Chem. Phys.)
12. "Experimental Second- and Third-order Molecular Hyperpolarizabilities," J.F. Ward and C.K. Miller. (Planned for submission to J. Chem. Phys.)
13. "Four-photon Parametric Oscillation in Cesium Vapor," A.V. Smith and J.F. Ward. (Planned.)

#### THESES PRESENTED

- K.M. Leung, "Some Resonant Processes in Nonlinear Optics," 1972.
- I.J. Bigio, "Studies in dc-Induced Optical Second-Harmonic Generation in Gases," 1973.

#### PAPERS PRESENTED

- Resonant Processes in Nonlinear Optics, J.F. Ward and K.M. Leung, Gordon Conference on Nonlinear Optics, Beaver Dam, Wisconsin, 23-28 July, 1972.
- Second- and Third-order Nonlinear Coefficients for Some Halogenated Methanes, I.J. Bigio, International Quantum Electronics Conference, San Francisco, June, 1974.
- Electric Field Induced Optical Rectification in Nitrobenzene, J.K. Guha and J.F. Ward, Bull. Am. Phys. Soc. 21, 99 (1976). Presented at 1976 Annual APS Meeting, New York, February, 1976.

#### SEMINARS GIVEN

J.F. Ward, "Resonant Nonlinear Optics," Colloquium at University of Toledo,  
1 Feb. 1973.

J.F. Ward, "Highlights of the Gordon Conference on Nonlinear Optics," U/M  
Physics Department Low Energy Seminar, Feb. 1974.

Irving J. Bigio, "Introduction to Nonlinear Optics," University of Michigan  
Student Colloquium, Jan. 1973.

Irving J. Bigio, "Second-Harmonic Generation in Gases," U/M Physics Department  
Low Energy Seminar, Oct. 1973.

Irving J. Bigio, "dc-Induced Second-Harmonic Generation in Gases," Lawrence  
Livermore Laboratory, Univ. of California, Oct. 1973.

Irving J. Bigio, "dc-Induced Second-Harmonic Generation in Gases," Center for  
Laser Studies, Univ. of Southern California, Oct. 1973.

Irving J. Bigio, "Second-Harmonic Generation in Gases," Esso Research and  
Engineering Co., Linden, New Jersey, Nov. 1973.

Irving J. Bigio, "Harmonic Generation in Gases," Sandia Laboratories, Albuquerque,  
New Mexico, Mar. 1974.

Irving J. Bigio, "Harmonic Generation in Gases," Los Alamos Scientific Labs,  
Los Alamos, New Mexico, Mar. 1974.

A. Smith, "Nonlinear Optics and Harmonic Generation in Cesium Vapor," U/M Physics  
Department Student Seminar, May 1974.

C.K. Miller, "Molecular Hyperpolarizabilities," Physical Chemistry Department,  
U/M, Jan. 1975.

J.F. Ward, "Resonant Nonlinear Optics," Los Alamos Scientific Lab, Jan. 1975.

J.F. Ward, "Resonant Nonlinear Optics," Optical Sciences Institute, University  
of Arizona, Jan. 1975.

J.F. Ward, "Resonant Nonlinear Optics," University of Southern California, Feb.  
1975.

J.F. Ward, "Resonant Nonlinear Optics," Seminar presented at Aerospace Corp.,  
Los Angeles, Apr. 1975.

J.K. Guha, "Spectroscopy and Laser Properties of Neodymium-Doped Laser Materials,"  
University of Toledo, Apr. 1975.

J.F. Ward, "Lasers and Nonlinear Optics," University of Michigan, Dearborn Campus

J.F. Ward, "Two Nonlinear Optical Processes: The Oldest and the Newest," Condensed  
Matter Seminar, U/M, Nov. 1975.

A.V. Smith, "Saturation in Optical Processes," Atomic Physics Seminar, U/M, Apr.  
1976.

C.K. Miller, "Optical Harmonic Generation in Gases," Research Seminar, U/M, Mar.  
1976.


#### PARTICIPANTS

J.F. Ward	Principal Investigator
K.M. Leung	Graduate Student and Research Associate, 1972. (Now at U.S.C.)
I.J. Bigio	Graduate Student 1972 - 1973. (Now at LASL.)
A.V. Smith	Graduate Student, 1972 -
C.K. Miller	Graduate Student, 1973 -
M. Myers	Graduate Student, 1976 -
D. Elliott	Graduate Student, working part time on the project.

The following have also participated on an hourly or unpaid basis:

E. Pellic, D. Kania.

Respectfully submitted,



J.F. Ward  
Professor of Physics  
University of Michigan



## APPENDIX

Molecule	$-\Delta k_0$ ( $\text{cm}^{-1}$ )	$10^{36} \chi_{\parallel}^{(3)}$ (esu)	$10^{30} \chi_{\parallel}^{(2)}$ (esu)
$\text{H}_2\text{S}$	$9.73 \pm .10$	$.865 \pm .022$	$-.043 \pm .009$
$\text{NH}_3$	$5.40 \pm .05$	$.512 \pm .010$	$-.210 \pm .004$
$(\text{CH}_3)_2\text{O}$	$8.92 \pm .09$	$.823 \pm .030$	$-.468 \pm .011$
$\text{CO}$	$3.35 \pm .03$	$.144 \pm .004$	$.129 \pm .013$
$\text{NO}$	$2.72 \pm .03$	$.235 \pm .007$	$.153 \pm .018$
$\text{H}_2\text{O}$	$2.42 \pm .02$	$.192 \pm .010$	$-.094 \pm .003$
$\text{H}_2$	$1.27 \pm .01$	$.0652 \pm .0008$	--
$\text{N}_2$	$1.89 \pm .02$	$.0866 \pm .0010$	--
$\text{O}_2$	$2.41 \pm .02$	$.0953 \pm .0016$	--
$\text{SF}_6$	$3.47 \pm .03$	$.130 \pm .002$	--
$\text{CO}_2$	$3.36 \pm .03$	$.1119 \pm .0013$	--